

# Metallic xenon. Polarizability. Equation of state.

V. N. Bogomolov

*Russian Academy of Sciences*

*A.F. Ioffe Physical & Technical Institute*

*St. Petersburg, Russia*

(February 1, 2008)

It is shown that some of the physical properties of inert gas ( IG ) condensates (polarizability, compressibility, metallization under pressure, equation of state) may be well described taking into account the first excited state of atoms only. The Herzfeld criterion of metallization well corresponds to the Mott transition criterion and to the percolation threshold. For metallic xenon concentration of the molecular type excitations corresponds to the Bose condensation temperature  $T_C \sim 4000K$ . The BCS formula gives  $T_c \sim 5000K$ . If phonons are changed by fluctuations of interatomic interaction energy. A simple relations between the parameters of atoms at metallization has been found.

One of the peculiarities of inert gas (IG) condensates is the coincidence of interatomic distances with diameters  $2r_2$  of atoms in the first excited state. This is the reason to ignore the next excited states. Such atoms are excimer analogies of the corresponding alkali metals. For this case an alternative approach to a description of properties of molecular condensates is possible [1]. These substances have a bond of a metallic type via excited state orbitals but without a metallic conductivity because a mean number of electrons at excited state orbitals  $X < 1$ . For xenon  $X = 0,038$ . Most of the physical properties ( condensation and adsorption energy, compressibility, metallization under pressure ) may be described by expressions of the theory of simple metals with electron charge  $e$  replaced by  $eX$  (and with replaced numeral coefficients) [2–4].

Properties of the condensate are determined by the excited state radius  $r_2$  of atoms. It may be expressed through energy by the hydrogen-like formula  $2r_2 = .e^2/(E_1 - E_2)$ . Here  $E_1 = e^2/2r_1$ — ionization potential,  $E_2$  - the transition energy between the ground and excited state of atom. For system in which the excited state radius of atoms is fixed the atomic wave functions may be expressed by the energy. Probability  $X$  for an electron to appear at excited state orbital with radius  $r_2$  may be obtained from the atomic wave functions of excited state:

$$X(r_2) = X_0 \exp \left[ -\frac{2E_2(r_2 - r_1)}{e^2} \right] = X_{01} \exp \left[ -\left( \frac{r_2}{r_1} + \frac{r_1}{r_2} \right) \right] = X_{02} \exp \left( -\frac{r_2}{r_1} \right) \text{ at } r_2 \gg r_1.$$

The pre-exponential coefficient is a relatively weak function of  $r$ . An average perturbation energy  $w = e^2/2(r_2 - r_1)$  is the interaction energy between electrons of the ground state orbitals of the neighbouring atoms at distance  $2r_2$  in condensate. The mean electron concentration  $X$  per atom depends on the perturbation energy  $w$ . The pressure is more significant parameter then the temperature. For ordinary semiconductors with well-defined band structure this energy is  $kT$ . For IG the relation is:  $g = w/E_2 \sim (0.30 - 0.55)$ . Metallization of xenon occurs at  $g_m = 0,75$ . The main properties of condensates may be expressed through atomic spectroscopic parameters of atoms. Radii of ground and excited states of atoms  $r_1$  and  $r_2$  may be changed by the corresponding atomic volumes  $V_1$  and  $V_2$ :

$$X(r_2) \sim \exp[-(y^{1/3} + y^{-1/3})]; \quad y = \frac{V_2}{V_1} = \frac{E_1^3}{(E_1 - E_2)^3}.$$

For IG volumes  $V_1$  are approximately independent on pressure:

|            | Ne    | Ar    | Kr    | Xe    |
|------------|-------|-------|-------|-------|
| $V_1(A^3)$ | 0.156 | 0.400 | 0.569 | 0.873 |

A simple physical picture of the model discussed may be revealed if applied to calculation of the polarizability of atoms in condensates.

Atomic polarizability  $a$  well corresponds to polarizability of the excited state orbital with the occupation number  $X$ :

$$a_c \sim eXr_2 \frac{er_2}{E_1 - E_2} \sim Xr_2^3 \sim XV_2 \sim (E_1 - E_2)^{-3} \exp \left[ -\frac{E_2^2}{E_1(E_1 - E_2)} \right].$$

The molar refractivity  $R$  is proportional to polarizability  $a$ . Hence the Herzfeld criterion of metallization [5] must be proportional to  $XV_2$ . We have:

|                  | Ne   | Ar   | Kr   | Xe   |
|------------------|------|------|------|------|
| $R(cm^3/mol)$    | 1.00 | 4.15 | 6.25 | 10.2 |
| $XV_2(cm^3/mol)$ | 1.57 | 4.49 | 6.60 | 10.2 |

From the other side, metallization occurs at  $X_p = 0, 12$  ( $y_m = 14.4$ ) as a transition over the percolation threshold [1,2]. Consequently the molar volume at metallization  $V_{02m} \sim V_{2m} = V_1 y_m = V_1 14.4 \sim E_1^{-3}$  for all IG. This relation well corresponds to criterion of the Mott transition too ( $V_{2m}/V_1 \sim 15$ ).

|                     |      |      |      |       |
|---------------------|------|------|------|-------|
| $V_{02m}(cm^3/mol)$ | 1.73 | 4.65 | 6.63 | 10.20 |
|---------------------|------|------|------|-------|

Quantum mechanics calculations gives [6]:

|                     |   |      |      |       |
|---------------------|---|------|------|-------|
| $V_{02m}(cm^3/mol)$ | — | 4.61 | 6.67 | 10.23 |
|---------------------|---|------|------|-------|

Metallic xenon contain virtual excimer  $Xe_2$  molecules in metal type lattice. The mean concentration of  $Xe_2$  is about  $0.12 \times 3.015 \cdot 10^{22}$  and the Bose condensation temperature  $T_C \sim 4000 K$  (at  $m^* = 2m_0$ ). By analogy with the BCS formula  $T_c \sim 0.5 w_m X_p \sim 0.5 E_2 X_p \sim 5000 K$  if phonons are changed by fluctuations of ineratomic interaction energy  $m$ . It does not contradict with [1]. For the hypothetical metallic  $H_2$  ( $H_4$  virtual molecules),  $T_C \sim 10000 K$  at  $V_{02m} \sim 3 cm^3/mol$ .

The next verification of the model may be made if used to calculation of the compressibility and equation of state (EOS). Metal-like compressibility:

$$-\frac{dV_2}{V_2 dP_c} \sim \frac{r_2^4}{e^2 X^2} \sim \frac{V^{4/3} y^{4/3}}{\exp[-2(y^{1/3} + y^{-1/3})]}$$

EOS at  $T = 0$  and  $y > 60$  is:

$$P_c(V_0) = P_0 V_1^{-4/3} \int (y-l)^{-7/3} \exp\left\{-2\left[(y-l)^{1/3} + (y-l)^{-1/3}\right]\right\} dy$$

$$= P_{10} E_1^4 F(E_1; V_0) + P_{20} = P_{30} F(E_1; V_0) + P_{20},$$

$$P_{10} = 173.74827(GPa; eV^{-7}); P_{20} = -1.294312(GPa);$$

$$P_{30} = 3.7578024 \cdot 10^6(GPa) \text{ at } E_1 = 12.127 eV \text{ for xenon.}$$

The volume  $V_1$  of atom in ground state is subtracted from the volume  $V_2$  (i.e. parameter  $y$  replaced by  $y - 1 = (V_2 - V_1)/V_1$ ).

The metallization pressure is  $P_m \sim E_1^4$  for molecular condensates.

The  $P_e$  vs.  $V_0$  data [7] are presented below together with the calculated  $P_c$  for xenon:

|                 |                   |       |      |       |       |       |       |       |       |      |       |
|-----------------|-------------------|-------|------|-------|-------|-------|-------|-------|-------|------|-------|
| $V_0(cm^3/mol)$ | 34.7              | 11.41 | 11.1 | 10.88 | 10.75 | 10.46 | 10.24 | 10.12 | 9.88  | 9.72 | 9.4   |
| $P_e(GPa)$      | $1 \cdot 10^{-4}$ | 107   | 117  | 125   | 130   | 142   | 152   | 158   | 171   | 180  | 200   |
| $P_c(GPa)$      | $1 \cdot 10^{-4}$ | 107.5 | 118  | 125.5 | 130   | 142   | 151.5 | 157   | 169.5 | 178  | 197.5 |
| $y$             | 48.80             | 16.1  | 15.6 | 15.3  | 15.1  | 14.7  | 14.4  | 14.2  | 13.9  | 13.7 | 13.2  |

Again, the agreement is good. The main purpose of these evaluations is to find extremely simple and adequate description of the intermolecular and metal-dielectric interactions alternative to traditional way. It may be useful for understanding of some problems of nanoelectronics and of multicomponent HTSC.

[1] Bogomolov V.N. Preprint <http://xxx.lanl.gov/abs/cond-mat/9902353>

[2] Bogomolov V.N. Tech. Phys. Lett. 1995. V.21. N11. P.928-930

[3] Bogomolov V.N. Phys. Sol. State 1993. V.35. N4. P.469-474

[4] Bogomolov V.N. Phys. Rev. 1995. V.51 N23. P.17040-17045

[5] Herzfeld K.F. Phys Rev. 1927. V. 29. N5. P.701-708

[6] Tulub A.V., Bratsev V.F., Pak M.V. Optika i Spectrosc., 1993. V.74, P.464-469

[7] Göttel K.A., Eggert J.H., Silvera I.F., Moss W.C. Phys. Rev. Lett. 1989. V. 62, N6, P.665-668